

Antibiotics: A Risk Factor for CDiff

High Risk for CDiff

- ☐ Clindamycin (Cleocin®)*
- ☐ Ceftriaxone (Rocephin®)*
- ☐ Ciprofloxacin (Cipro®)*
- ☐ Levofloxacin (Levaquin®)*
- ☐ Cefepime (Maxipime®)
- ☐ Ceftazidime (Fortaz®)
- ☐ Cefuroxime (Ceftin®)
- ☐ Ertapenem (Invanz®)
- ☐ Meropenem (Merrem®)

Medium Risk for CDiff

- ☐ Piperacillin/tazobactam (Zosyn®)*
- ☐ Amoxicillin/clavulanic acid (Augmentin®)*
- ☐ Ampicillin/sulbactam (Unasyn®)
- ☐ Amoxicillin (Amoxil®)
- ☐ Ampicillin
- ☐ Azithromax (Zithromax®)
- ☐ Aztreonam (Azactam®)
- ☐ Cefazolin (Ancef®)
- ☐ Cephalexin (Keflex®)
- ☐ Dalfopristin/-quinupristin (Synercid®)

Low Risk for CDiff

- ☐ Amikacin (Amikin®)
- ☐ Daptomycin (Cubicin®)
- ☐ Doxycycline (Vibramycin®)
- ☐ Fosfomycin (Monurol®)
- ☐ Gentamicin
- ☐ Linezolid (Zyvox®)
- ☐ Nitrofurantoin (Macrobid®)
- ☐ Polymixin (Colistin®)
- ☐ Rifampin (Rifadin®)
- ☐ Trimethoprim/-sulfamethoxazole (Bactrim®)

*Highest Association with CDiff